



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

**Altered monoaminergic systems and depressive-like behavior in congenic
prion protein nnock-out mice**

Nuvolone, Mario ; Aguzzi, Adriano

DOI: <https://doi.org/10.1074/jbc.L115.689117>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-114215>

Journal Article

Published Version

Originally published at:

Nuvolone, Mario; Aguzzi, Adriano (2015). Altered monoaminergic systems and depressive-like behavior in congenic prion protein nnock-out mice. *Journal of Biological Chemistry*, 290(43):26350.

DOI: <https://doi.org/10.1074/jbc.L115.689117>

LETTER

Altered Monoaminergic Systems and Depressive-like Behavior in Congenic Prion Protein Knock-out Mice

Beckman *et al.* (1) reported altered monoaminergic systems and depressive-like behaviors in a line of mice lacking the cellular prion protein PrP^C (congenic B10.129-*Prnp*^{Edbg/Edbg} mice). These mice were derived from gene-targeted 129/Ola embryonic stem cells and were subsequently crossed to C57BL/10SnJ mice. Consequently, the genomic region of B10.129-*Prnp*^{Edbg/Edbg} mice flanking the targeted *Prnp* locus contains a stretch of up to 47.4 Mb isogenic to 129/Ola (2). In this region, any genes polymorphic between the 129/Ola and C57BL/10SnJ strains will display 129/Ola allelotypes (2). This genetic anomaly represents a systematic confounder when congenic knock-out mice are compared with wild-type mice of the back-crossing strain, even when using littermates from heterozygous breedings (2). This phenomenon, intrinsic to all congenic knock-out mice, is known as the flanking gene problem.

The genetic background has a strong influence on depression-related behavior in mice (3). Numerous studies have identified multiple genes and quantitative trait loci controlling monoaminergic systems and depressive-like behaviors. Crucially, these include *Sirpa*, which is linked to *Prnp* (genetic distance: approximately 2 centimorgans), is

polymorphic between C57BL and 129 strains, and is a known genetic confounder of studies with *Prnp*^{-/-} mice (4, 5).

For the above reasons, the published data (1) do not allow conclusive determination of whether the phenotypes observed are caused by the ablation of *Prnp*, by the *Sirpa* polymorphism discussed above, or by other sequence polymorphisms in the genomic region flanking *Prnp*. The conclusion that PrP^C participates in the modulation of monoaminergic systems and depressive-like behavior is therefore unwarranted.

Mario Nuvolone¹ and Adriano Aguzzi¹

Institute of Neuropathology, University Hospital of Zurich, Zurich, Switzerland

1. Beckman, D., Santos, L. E., Americo, T. A., Ledo, J. H., de Mello, F. G., and Linden, R. (2015) Prion protein modulates monoaminergic systems and depressive-like behavior in mice. *J. Biol. Chem.* **290**, 20488–20498
2. Striebel, J. F., Race, B., and Chesebro, B. (2013) Prion protein and susceptibility to kainate-induced seizures: genetic pitfalls in the use of PrP knockout mice. *Prion* **7**, 280–285
3. Jacobson, L. H., and Cryan, J. F. (2007) Feeling strained? Influence of genetic background on depression-related behavior in mice: a review. *Behav. Genet.* **37**, 171–213
4. Nuvolone, M., Kana, V., Hutter, G., Sakata, D., Mortin-Toth, S. M., Russo, G., Danska, J. S., and Aguzzi, A. (2013) SIRPα polymorphisms, but not the prion protein, control phagocytosis of apoptotic cells. *J. Exp. Med.* **210**, 2539–2552
5. Ohnishi, H., Murata, T., Kusakari, S., Hayashi, Y., Takao, K., Maruyama, T., Ago, Y., Koda, K., Jin, F. J., Okawa, K., Oldenborg, P. A., Okazawa, H., Murata, Y., Furuya, N., Matsuda, T., Miyakawa, T., and Matozaki, T. (2010) Stress-evoked tyrosine phosphorylation of signal regulatory protein α regulates behavioral immobility in the forced swim test. *J. Neurosci.* **30**, 10472–10483

DOI 10.1074/jbc.L115.689117

¹E-mail: adriano.aguzzi@usz.ch or mario.nuvolone@usz.ch

Altered Monoaminergic Systems and Depressive-like Behavior in Congenic Prion Protein Knock-out Mice

Mario Nuvolone and Adriano Aguzzi

J. Biol. Chem. 2015, 290:26350.

doi: 10.1074/jbc.L115.689117

Access the most updated version of this article at <http://www.jbc.org/content/290/43/26350>

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 5 references, 3 of which can be accessed free at <http://www.jbc.org/content/290/43/26350.full.html#ref-list-1>